



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

tically all nucleus. The corpuscles on the other hand have lost their nuclei wholly. Between these extremes we have various stages of chromatin reduction in the development of the specialized Metazoan tissues. The maturation divisions in ova and sperm, the bodily extrusion of chromatin observed on the part of blood-cells, etc., he regards as illustrations of the process.

#### THE RESERVE OF FOOD IN TREES

Preston and Phillips (Forest Quart., 1911) agree with the common view that starch is the principal form in which reserve food is stored in trees. They doubt that cellulose is able to act at all as a reserve material. The maximum contained reserve for deciduous trees occurs about the time the leaves fall, and during the next few weeks there is a decided reduction in its amount. The sugar content in trees remains pretty constant except for an increase in spring during the unfolding of the buds.

#### ALTERNATION OF GENERATION IN FLORIDEÆ

Lewis (Bot. Gaz., Mch., 1912), by artificial plantings of tetraspores and carpospores of *Polysiphonia* and some other genera of red algae gets experimental results supporting the general conclusion that tetraspores produce only the sexual plants and carpospores only the tetrasporic plants. In no instance was an exception found to the rule, although a considerable number of plantings developed to maturity. Tetraspores from a given individual produced male and female plants in approximately equal numbers. It is also concluded that no greater growth vigor comes to the carpospores over the tetraspores because of the double number of chromosomes contained by them.

#### RELATION OF THE PROTOPLASM OF ADJACENT PROTOPLASTS

Thoday (Ann. Bot., 1911) undertakes to throw light on the relation that exists between protoplasts of contiguous cells, by an examination of the relation between the parasite, *Cuscuta*, and its host. She finds that there is no direct protoplasmic connection between the cells of *Cuscuta* and the host, but that the phloem cells of the parasite haustoria apply themselves to the sieve plates of the phloem of

the host. The wall of the former breaks down over the area of contact, but the protoplasmic threads do not penetrate and become continuous with the protoplasm of the sieve cells. The parasite seems to receive the food passively through the action of internal pressure in the cells of the host.

The author suggests that this study appears to lend support to Gardiner's view that connecting threads of protoplasm, as found between the cells of some plants, represent original connections between genetically connected cells due to lack of complete dividing walls along the plane of division, and are not secondary connections of the protoplasts effected by the cell walls being perforated by advancing protoplasmic fibrils.

#### A NEW FORM OF PARASITISM

Jensen (Ann. Jard. Bot. Buitenzorg, 1910) reports the study of several species of dipterous larvae, belonging to not less than three different families, that are able to live in the mixture found in pitchers of *Nepenthes*. They are able to resist the action of the ferments secreted by the *Nepenthes*, which digests the other contained organisms. He found that closely related larvae taken from nearby waters were wholly unable to withstand the action of the fluid. These conditions seem quite analogous to those encountered and overcome by some intestinal parasites of animals.

#### BACTERIA IN RIVER WATER

Reiss (Verh. Phys. Med. Gessell. Wurtzburg, 1911) reports the finding of as many as sixty-two species of bacteria in samples of river water taken from the Main, near Wurtzburg. On cultivation, a number of species showed resemblances to known pathogenic types, but seemed to have lost their virulence. This is certainly a remarkable showing of bacteria.

#### CONDITION OF CONTINUOUS DIVISION IN PARAMECIUM

Woodruff (Biol. Bull., Dec., 1911) believes that most, if not all, normal individuals of *Paramecium*, under suitable environmental conditions, possess unlimited power of reproduction without conjugation or artificial stimulation; and that the discrepant results of dif-